

Spacelike Weingarten Surfaces in R_1^3 and the Sine-Gordon Equation*

Chen Weihuan[†]

Department of Mathematics, Peking University, Beijing, 100871.

metadata, citation and similar papers at core.ac.uk

and

Li Haizhong[‡]

*Department of Applied Mathematics, Tsinghua University, Beijing, 100084,
People's Republic of China*

Submitted by Colin Rogers

Received February 5, 1997

In this paper we will establish a relation between spacelike Weingarten surfaces with condition $K + 2mH = m^2 + l^2$ in R_1^3 and solutions of the sine-Gordon equation. Moreover, we will give a method to construct new spacelike Weingarten surfaces from a given spacelike Weingarten surface in R_1^3 by Bäcklund transformation. © 1997 Academic Press

1. SPACELIKE WEINGARTEN SURFACES

Let R_1^3 be a 3-dimensional Minkowski space. We consider a spacelike Weingarten surface S in R_1^3 satisfying

$$K + 2mH = m^2 + l^2, \quad (1.1)$$

in which K is its Gauss curvature, H the mean curvature, and m, l are constants with $m^2 + l^2 > 0$.

* This work is supported in part by the NNSF of China.

[†] E-mail address: whchen@sxx0.math.pku.edu.cn.

[‡] E-mail address: hli@math.tsinghua.edu.cn.

Suppose that S is a spacelike surface with condition (1.1) in R_1^3 , and S has no umbilical point, so we can take the lines of curvature as its parametric curves, and write its fundamental forms as

$$\begin{aligned} I &= A^2 du^2 + B^2 dv^2, \\ II &= k_1 A^2 du^2 + k_2 B^2 dv^2, \end{aligned}$$

where k_1, k_2 are the principal curvatures of S . Let

$$\begin{aligned} \omega^1 &= A du, & \omega^2 &= B dv, \\ \omega_1^3 &= k_1 A du, & \omega_2^3 &= k_2 B dv. \end{aligned} \quad (1.2)$$

Then the Gauss and Codazzi equations become

$$\begin{aligned} K &= -k_1 k_2, \\ (k_1 - k_2) A_v + (k_1)_v A &= 0, \\ (k_2 - k_1) B_u + (k_2)_u B &= 0. \end{aligned} \quad (1.3)$$

By use of the same method as in our previous paper [2], we can introduce new parameters locally on S , denoted still by u and v , such that its fundamental forms become

$$\begin{aligned} I &= \frac{1}{m^2 + l^2} \left(\cos^2 \frac{\phi}{2} \cdot du^2 + \sin^2 \frac{\phi}{2} \cdot dv^2 \right), \\ II &= \frac{1}{\sqrt{m^2 + l^2}} \left(\sin \frac{\phi - \phi_0}{2} \cos \frac{\phi}{2} \cdot du^2 - \cos \frac{\phi - \phi_0}{2} \sin \frac{\phi}{2} \cdot dv^2 \right), \end{aligned} \quad (1.4)$$

where

$$\sin \frac{\phi_0}{2} = -\frac{m}{\sqrt{m^2 + l^2}}, \quad \cos \frac{\phi_0}{2} = \frac{l}{\sqrt{m^2 + l^2}}. \quad (1.5)$$

For convenience, the local parameters u, v of a spacelike Weingarten surface S with condition (1.1), for which its fundamental forms can be written by (1.4), are called *Tschebyscheff coordinates* on S , and ϕ is called the *Tschebyscheff angle*.

It follows from (1.4) that

$$\omega^1 = \frac{1}{\sqrt{m^2 + l^2}} \cos \frac{\phi}{2} \cdot du, \quad \omega^2 = \frac{1}{\sqrt{m^2 + l^2}} \sin \frac{\phi}{2} \cdot dv, \quad (1.6)$$

$$\begin{aligned}\omega_1^3 &= \omega_3^1 = -k_1 \omega^1 = -\sin \frac{\phi - \phi_0}{2} \cdot du, \\ \omega_2^3 &= \omega_3^2 = -k_2 \omega^2 = \cos \frac{\phi - \phi_0}{2} \cdot dv,\end{aligned}\tag{1.7}$$

and

$$\omega_1^2 = -\omega_2^1 = \frac{1}{2}(\phi_v du + \phi_u dv).\tag{1.8}$$

The principal curvatures of S are

$$\begin{aligned}k_1 &= \sqrt{m^2 + l^2} \cdot \frac{\sin((\phi - \phi_0)/2)}{\cos(\phi/2)}, \\ k_2 &= -\sqrt{m^2 + l^2} \cdot \frac{\cos((\phi - \phi_0)/2)}{\sin(\phi/2)}.\end{aligned}\tag{1.9}$$

Obviously, the following Codazzi equations

$$d\omega_1^3 = \omega_1^2 \wedge \omega_2^3, \quad d\omega_2^3 = \omega_2^1 \wedge \omega_1^3$$

hold automatically and the Gauss equation $d\omega_1^2 = K\omega^1 \wedge \omega^2 = \omega_1^3 \wedge \omega_2^3$ reads

$$\phi_{uu} - \phi_{vv} = -\sin(\phi - \phi_0).\tag{1.10}$$

We record these results as follows.

THEOREM 1.1. *Suppose S is a spacelike Weingarten surface without umbilics in R_1^3 satisfying condition (1.1). Then its Tschebyscheff angle ϕ is a solution of the sine-Gordon equation (1.10).*

Conversely, if $\tilde{\phi}$ is a solution of the sine-Gordon equation

$$\tilde{\phi}_{uu} - \tilde{\phi}_{vv} = -\sin \tilde{\phi},\tag{1.11}$$

and m, l are constants such that $m^2 + l^2 > 0$, then there exists locally a spacelike Weingarten surface S with condition (1.1) in R_1^3 such that $\phi = \tilde{\phi} + \phi_0$ is its Tschebyscheff angle, where ϕ_0 is given by (1.5).

2. DARBOUX LINE CONGRUENCE IN R_1^3

DEFINITION 2.1. Suppose S and S^* are two spacelike surfaces in R_1^3 , and there exists a correspondence between them such that

(1) $\overrightarrow{PP^*}$ the distance between corresponding points is a constant λ , that is, $|\overrightarrow{PP^*}| = \lambda$, where $|v| = \sqrt{v \cdot v}$ for a spacelike vector v , and $|v| = \sqrt{-v \cdot v}$ for a timelike vector v ;

(2) lines joining corresponding points are isoclinic with the surfaces S and S^* , that is, we can choose unit normal vectors e_3 and e_3^* on S and S^* , respectively, such that $t \cdot e_3 = -t \cdot e_3^* = \text{constant}$, where t is the unit direction vector of $\overrightarrow{PP^*}$;

(3) the inner product of unit normal vectors at corresponding points of S and S^* is constant, i.e., $e_3 \cdot e_3^* = \text{constant}$.

Then the congruence in R_1^3 , formed by lines joining corresponding points of S and S^* , is called a Darboux line congruence associated with them.

Remark 2.1. Evidently, a pseudo-spherical line congruence between spacelike surfaces in R_1^3 (see [8]) is a special kind of Darboux line congruence (i.e., $t \cdot e_3 = 0 = t \cdot e_3^*$).

A Darboux line congruence is called spacelike or timelike according to its direction vector t being spacelike or timelike (the case that t is lightlike is not in our consideration).

If Σ is a timelike Darboux line congruence, then we can assume that

$$t \cdot e_3 = -t \cdot e_3^* = -\cosh \gamma = \text{const.}, \quad (2.1)$$

$$e_3 \cdot e_3^* = \cosh \theta = \text{const.}, \quad (2.2)$$

in this case, the end points of vectors $t, e_3, -e_3^*$ lie on the hyperbolic space

$$H^2 = \{(x^1, x^2, x^3) \in R_1^3 : (x^1)^2 + (x^2)^2 - (x^3)^2 = -1, x^3 > 0\},$$

where γ is the geodesic distance between t and e_3 , and θ is the geodesic distance between e_3 and e_3^* . By the triangle inequality, we have

$$2\gamma \geq \theta > 0,$$

and

$$\coth \gamma \leq \coth \frac{\theta}{2},$$

i.e.,

$$\sinh^2 \theta - (\cosh \theta - 1)^2 \coth^2 \gamma \geq 0. \quad (2.3)$$

If Σ is a spacelike Darboux line congruence, we can assume that

$$t \cdot e_3 = -t \cdot e_3^* = -\sinh \gamma = \text{const.},$$

$$e_3 \cdot e_3^* = \cosh \theta = \text{const.},$$

then we always have

$$\sinh^2 \theta - (\cosh \theta - 1)^2 \tanh^2 \gamma = \sinh^2 \theta \left(1 - \tanh^2 \frac{\theta}{2} \tanh^2 \gamma \right) \geq 0. \quad (2.4)$$

THEOREM 2.1. *Suppose that S and S^* are two spacelike surfaces in R_1^3 , and there exists a correspondence between them such that the lines joining corresponding points form a Darboux line congruence Σ in R_1^3 . Then both S and S^* are spacelike Weingarten surfaces satisfying the same condition as (1.1), in which*

(1) *when Σ is a timelike Darboux line congruence,*

$$m = \frac{(\cosh \theta - 1) \cosh \gamma}{\lambda \sinh^2 \gamma}, \quad l^2 = \frac{\sinh^2 \theta}{\lambda^2 \sinh^2 \gamma} \left(1 - \tanh^2 \frac{\theta}{2} \cdot \coth^2 \gamma \right), \quad (2.5)$$

(2) *when Σ is a spacelike Darboux line congruence,*

$$m = \frac{(\cosh \theta - 1) \sinh \gamma}{\lambda \cosh^2 \gamma}, \quad l^2 = \frac{\sinh^2 \theta}{\lambda^2 \cosh^2 \gamma} \left(1 - \tanh^2 \frac{\theta}{2} \cdot \tanh^2 \gamma \right). \quad (2.6)$$

Proof. Case (1). Let Σ be a timelike Darboux line congruence associated with spacelike surfaces S and S^* . Then for a local orthonormal frame field $\{r; e_1, e_2, e_3\}$ on S , the direction vector t along congruence Σ can be expressed as

$$t = \sinh \gamma (\cos \psi e_1 + \sin \psi e_2) + \cosh \gamma e_3. \quad (2.7)$$

So the position vector r^* of S^* can be expressed as

$$r^* = r + \lambda t, \quad (2.8)$$

where $\lambda > 0$ is constant. We can choose the normal vector e_3^* of S^* so that

$$e_3 \cdot e_3^* = \cosh \theta, \quad t \cdot e_3 = -t \cdot e_3^* = -\cosh \gamma, \quad e_3^* \cdot e_3^* = -1, \quad (2.9)$$

where both θ and γ are constant. Now, the normal vector e_3^* of S^* can be also written as

$$e_3^* = x_1 e_1 + x_2 e_2 - \cosh \theta e_3. \quad (2.10)$$

Then we get from (2.9)

$$\begin{aligned}x_1 &= -(\cosh \theta - 1) \coth \gamma \cos \psi \\&\quad + \sqrt{\sinh^2 \theta - (\cosh \theta - 1)^2 \coth^2 \gamma} \cdot \sin \psi, \\x_2 &= -(\cosh \theta - 1) \coth \gamma \sin \psi \\&\quad - \sqrt{\sinh^2 \theta - (\cosh \theta - 1)^2 \coth^2 \gamma} \cdot \cos \psi.\end{aligned}$$

Put

$$\begin{aligned}\sin \frac{\phi_0}{2} &= -\frac{\cosh \theta - 1}{\sinh \theta} \coth \gamma, \\ \cos \frac{\phi_0}{2} &= \frac{\sqrt{\sinh^2 \theta - (\cosh \theta - 1)^2 \coth^2 \gamma}}{\sinh \theta}.\end{aligned}\tag{2.11}$$

Then

$$x_1 = \sinh \theta \sin \left(\psi + \frac{\phi_0}{2} \right), \quad x_2 = -\sinh \theta \cos \left(\psi + \frac{\phi_0}{2} \right). \tag{2.12}$$

Taking differentiation of (2.8) and using the structure equations we get

$$\begin{aligned}dr^* &= (\omega_1 + \lambda \cosh \gamma \omega_1^3) e_1 + (\omega^2 + \lambda \cosh \gamma \omega_2^3) e_2 \\&\quad + \lambda \sinh \gamma (d\psi + \omega_1^2) (-\sin \psi e_1 + \cos \psi e_2) \\&\quad + \lambda \sinh \gamma (\cos \psi \omega_1^3 + \sin \psi \omega_2^3) e_3.\end{aligned}\tag{2.13}$$

From $dr^* \cdot e_3^* = 0$ we have

$$\begin{aligned}&\lambda \sinh \gamma \sinh \theta \cos \frac{\phi_0}{2} (d\psi + \omega_1^2) \\&= \sinh \theta \left(\sin \left(\psi + \frac{\phi_0}{2} \right) \omega^1 - \cos \left(\psi + \frac{\phi_0}{2} \right) \omega^2 \right) \\&\quad + \lambda \sinh \theta \cosh \gamma \left(\sin \left(\psi + \frac{\phi_0}{2} \right) \omega_1^3 \right. \\&\quad \left. - \cos \left(\psi + \frac{\phi_0}{2} \right) \omega_2^3 \right) + \lambda \cosh \theta \sinh \gamma (\cos \psi \omega_1^3 + \sin \psi \omega_2^3). \tag{2.14}\end{aligned}$$

Taking differentiation of (2.14) and using the structure equation we get

$$\begin{aligned} & \lambda \sinh \gamma \sinh \theta \cos \frac{\phi_0}{2} \omega_1^3 \wedge \omega_2^3 \\ &= (d\psi + \omega_1^2) \wedge \left\{ \sinh \theta \left(\cos \left(\psi + \frac{\psi_0}{2} \right) \omega^1 + \sin \left(\phi + \frac{\phi_0}{2} \right) \omega^2 \right) \right. \\ & \quad + \lambda \sinh \theta \cosh \gamma \left(\cos \left(\psi + \frac{\phi_0}{2} \right) \omega_1^3 + \sin \left(\psi + \frac{\phi_0}{2} \right) \omega_2^3 \right) \\ & \quad \left. + \lambda \cosh \theta \sinh \gamma (-\sin \psi \omega_1^3 + \cos \psi \omega_2^3) \right\}. \end{aligned}$$

Then by putting (2.14) in it we have

$$\begin{aligned} & \lambda^2 \sinh^2 \gamma \sinh^2 \theta \cos^2 \frac{\phi_0}{2} \omega_1^3 \wedge \omega_2^3 \\ &= -\sinh^2 \theta \omega^1 \wedge \omega^2 \\ & \quad - \lambda \left(\sinh^2 \theta \cosh \gamma + \sinh \theta \cosh \theta \sinh \gamma \sin \frac{\phi_0}{2} \right) \\ & \quad \times (\omega^1 \wedge \omega_2^3 + \omega_1^3 \wedge \omega^2) \\ & \quad - \lambda^2 \left(\sinh^2 \theta \cosh^2 \gamma + \cosh^2 \theta \sinh^2 \gamma \right. \\ & \quad \left. + 2 \sinh \theta \cosh \theta \sinh \gamma \cosh \gamma \sin \frac{\phi_0}{2} \right) \omega_1^3 \wedge \omega_2^3. \quad (2.15) \end{aligned}$$

We know that

$$\begin{aligned} \omega^1 \wedge \omega_2^3 + \omega_1^3 \wedge \omega^2 &= -(k_1 + k_2) \omega^1 \wedge \omega^2, \\ \omega_1^3 \wedge \omega_2^3 &= k_1 k_2 \cdot \omega^1 \wedge \omega^2. \end{aligned} \quad (2.16)$$

Then (2.14) can be reduced to

$$k_1 k_2 - \frac{(\cosh \theta - 1) \cosh \gamma}{\lambda \sinh^2 \gamma} (k_1 + k_2) + \frac{\sinh^2 \theta}{\lambda^2 \sinh^2 \gamma} = 0. \quad (2.17)$$

By comparing (2.17) with (1.1) we obtain (2.5).

Because (2.8) can be written as

$$r = r^* + \lambda(-t),$$

the same argument is valid for S^* . Then the surface S^* satisfies the same condition (2.17).

Case (2). Let Σ be a spacelike Darboux line congruence associated with spacelike surfaces S and S^* . Then for a local orthogonal frame field $\{r; e_1, e_2, e_3\}$ on S , the direction vector t along congruence Σ can be expressed as

$$t = \cosh \gamma (\cos \psi e_1 + \sin \psi e_2) + \sinh \gamma e_3. \quad (2.18)$$

By a similar procedure as in Case (1) we obtain

$$\begin{aligned} & \lambda \cosh \gamma \sinh \theta \cos \frac{\phi_0}{2} (d\psi + \omega_1^2) \\ &= \sinh \theta \left(\sin \left(\psi + \frac{\phi_0}{2} \right) \omega^1 - \cos \left(\psi + \frac{\phi_0}{2} \right) \omega^2 \right) \\ & \quad + \lambda \sinh \theta \sinh \gamma \left(\sin \left(\psi + \frac{\phi_0}{2} \right) \omega_1^3 \right. \\ & \quad \left. - \cos \left(\psi + \frac{\phi_0}{2} \right) \omega_2^3 \right) + \lambda \cosh \theta \cosh \gamma (\cos \psi \omega_1^3 + \sin \psi \omega_2^3), \end{aligned} \quad (2.19)$$

in which ϕ_0 is given by

$$\begin{aligned} \sin \frac{\phi_0}{2} &= -\frac{\cosh \theta - 1}{\sinh \theta} \cdot \tanh \gamma, \\ \cos \frac{\phi_0}{2} &= \frac{\sqrt{\sinh^2 \theta - (\cosh \theta - 1)^2 \tanh^2 \gamma}}{\sinh \theta}. \end{aligned} \quad (2.20)$$

A similar argument as above gives

$$k_1 k_2 - \frac{(\cosh \theta - 1) \sinh \gamma}{\lambda \cosh^2 \gamma} (k_1 + k_2) + \frac{\sinh^2 \theta}{\lambda^2 \cosh^2 \gamma} = 0. \quad (2.21)$$

3. BÄCKLUND TRANSFORMATION

Suppose Σ is a timelike Darboux line congruence associated with spacelike surfaces S and S^* in R_1^3 , so S and S^* are Weingarten spacelike surfaces with condition (1.1), and m, l are given by (2.5). Let u, v be the

Tschebyscheff coordinates on S . Then

$$\begin{aligned}\omega^1 &= \frac{\lambda \sinh \gamma}{\sinh \theta} \cos \frac{\phi}{2} \cdot du, & \omega^2 &= \frac{\lambda \sinh \gamma}{\sinh \theta} \sin \frac{\phi}{2} \cdot dv, \\ \omega_1^3 &= -\sin \frac{\phi - \phi_0}{2} \cdot du, & \omega_2^3 &= \cos \frac{\phi - \phi_0}{2} \cdot dv, \\ \omega_1^2 &= \frac{1}{2}(\phi_v du + \phi_u dv),\end{aligned}\tag{3.1}$$

where ϕ_0 is given by (2.11). Let

$$\psi = -\frac{\phi^*}{2},\tag{3.2}$$

and put (3.1) into (2.14). Then we have

$$\begin{aligned}\sinh \theta (d\psi + \omega_1^2) &= -du \cdot \left(\sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} \right. \\ &\quad \left. + \cosh \theta \cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} \right) \\ &\quad - dv \cdot \left(\cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} \right. \\ &\quad \left. + \cosh \theta \sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} \right),\end{aligned}\tag{3.3}$$

that is,

$$\begin{aligned}\sinh \theta \cdot \frac{\phi_u^* - \phi_v}{2} &= \sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} \\ &\quad + \cosh \theta \cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2}, \\ \sinh \theta \cdot \frac{\phi_v^* - \phi_u}{2} &= \cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} \\ &\quad + \cosh \theta \sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2}.\end{aligned}\tag{3.4}$$

By verifying directly we get the following

PROPOSITION 3.1. *If ϕ is a solution of the sine-Gordon equation*

$$\phi_{uu} - \phi_{vv} = -\sin(\phi - \phi_0), \quad (3.5)$$

then, for any $\theta \neq 0$, Eqs. (3.4) are completely integrable for ϕ^ , and its solution ϕ^* satisfies the same Eq. (3.5).*

Therefore (3.4) give a Bäcklund transformation of solutions of the sine-Gordon equation (3.5). Furthermore, if Σ is a spacelike Darboux line congruence associated with spacelike surfaces S and S^* in R_1^3 , then we obtain the same Bäcklund transformation (3.4) from (2.19).

In fact, we can prove the following theorem.

THEOREM 3.1. *Suppose Σ is a spacelike (or timelike) Darboux line congruence associated with spacelike surfaces in R_1^3 . Then both S and S^* are spacelike Weingarten surfaces satisfying the same condition as (1.1), and their Tscheytscheff angles ϕ and ϕ^* are both solutions of the sine-Gordon equation (3.5), and they are related to each other by a Bäcklund transformation (3.4), where ϕ_0 is given by (1.5).*

Proof. We will prove this theorem only for Σ being the timelike case, and the proof in another case is similar.

Recalling the definition of cross product of vectors in R_1^3 (see [1]), we have

$$\begin{aligned} e_3 \times e_3^* &= x^1 \cdot e_3 \times e_1 + x^2 \cdot e_3 \times e_2 = x^2 e_1 - x^1 e_2 \\ &= -\sinh \theta \left(\cos \left(\psi + \frac{\phi_0}{2} \right) e_1 + \sin \left(\psi + \frac{\phi_0}{2} \right) e_2 \right) \\ &= -\sinh \theta \left(\cos \frac{\phi_0}{2} \cdot \tau + \sin \frac{\phi_0}{2} \cdot \tau^\perp \right), \end{aligned} \quad (3.6)$$

where

$$\tau = \cos \psi \cdot e_1 + \sin \psi \cdot e_2 \quad (3.7)$$

is the unit vector of orthogonal projection of the direction vector t of Σ on tangent space of S , and

$$\tau^\perp = -\sin \psi \cdot e_1 + \cos \psi \cdot e_2.$$

Then the vector

$$e = \cos \frac{\phi_0}{2} \cdot \tau + \sin \frac{\phi_0}{2} \cdot \tau^\perp = \cos \left(\psi + \frac{\phi_0}{2} \right) e_1 + \sin \left(\psi + \frac{\phi_0}{2} \right) e_2 \quad (3.8)$$

is tangent to both S and S^* , and $\phi_0/2$ is the angle between τ and the vector e .

As stated at the beginning of this section we have a local frame field $\{r; e_1, e_2, e_3\}$, in which e_1, e_2 are along the principle directions of S . In addition we have two other local frame fields $\{r; \tau, \tau^\perp, e_3\}$ and $\{r; e, e^\perp, e_3\}$ on S , where

$$\begin{aligned} e^\perp &= e \times e_3 = -\sin \frac{\phi_0}{2} \cdot \tau + \cos \frac{\phi_0}{2} \cdot \tau^\perp \\ &= -\sin \left(\psi + \frac{\phi_0}{2} \right) e_1 + \cos \left(\psi + \frac{\phi_0}{2} \right) e_2, \end{aligned} \quad (3.9)$$

and a local frame field $\{r^*; e^*, e^{*\perp}, e_3^*\}$ on S^* , where

$$\begin{aligned} e^* &= -e = -\cos \left(\psi^* - \frac{\phi_0}{2} \right) \cdot e_1 - \sin \left(\psi^* - \frac{\phi_0}{2} \right) \cdot e_2 \\ &= -\cos \frac{\phi^* - \phi_0}{2} \cdot e_1 + \sin \frac{\phi^* - \phi_0}{2} \cdot e_2, \\ e^{*\perp} &= e^* \times e_3^* \\ &= \cosh \theta \left(-\sin \left(\psi + \frac{\phi_0}{2} \right) \cdot e_1 + \cos \left(\psi + \frac{\phi_0}{2} \right) \cdot e_2 \right) + \sinh \theta \cdot e_3 \\ &= \cosh \theta \left(\sin \frac{\phi^* - \phi_0}{2} \cdot e_1 + \cos \frac{\phi^* - \phi_0}{2} \cdot e_2 \right) + \sinh \theta \cdot e_3. \end{aligned} \quad (3.10)$$

Putting (3.3) into (2.13), we get

$$\begin{aligned} dr^* &= \left(\frac{\lambda \sinh \gamma}{\sinh \theta} \cos \frac{\phi}{2} - \lambda \cosh \gamma \sin \frac{\phi - \phi_0}{2} \right) du \cdot e_1 \\ &\quad + \left(\frac{\lambda \sinh \gamma}{\sinh \theta} \sin \frac{\phi}{2} + \lambda \cosh \gamma \cos \frac{\phi - \phi_0}{2} \right) dv \cdot e_2 \\ &\quad - \frac{\lambda \sinh \gamma}{\sinh \theta} \left\{ \left(\sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} \right. \right. \end{aligned}$$

$$\begin{aligned}
& + \cosh \theta \cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} \Big) du \\
& + \left(\cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} \right. \\
& \quad \left. + \cosh \theta \sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} \right) dv \Big\} \cdot \tau^\perp \\
& - \lambda \sinh \gamma \left(\cos \frac{\phi^*}{2} \sin \frac{\phi - \phi_0}{2} du + \sin \frac{\phi^*}{2} \cos \frac{\phi - \phi_0}{2} dv \right) \cdot e_3.
\end{aligned} \tag{3.11}$$

Denote $\{\xi^{*1}, \xi^{*2}\}$ the coframe dual to $\{r^*, e^*, e^{*\perp}, e_3^*\}$ on S^* . So

$$\begin{aligned}
\xi^{*1} &= dr^* \cdot e^* \\
&= -\frac{\lambda \sinh \gamma}{\sinh \theta} \left(\cos \frac{\phi^*}{2} \cos \frac{\phi - \phi_0}{2} du - \sin \frac{\phi^*}{2} \sin \frac{\phi - \phi_0}{2} dv \right), \\
\xi^{*2} &= dr^* \cdot e^{*\perp} \\
&= -\frac{\lambda \sinh \gamma}{\sinh \theta} \left(\cos \frac{\phi^*}{2} \sin \frac{\phi - \phi_0}{2} du + \sin \frac{\phi^*}{2} \cos \frac{\phi - \phi_0}{2} dv \right).
\end{aligned} \tag{3.12}$$

To find out ξ_1^{*3} and ξ_2^{*3} we take differentiation of (2.10), and get

$$\begin{aligned}
de_3^* &= (-x^2 e_1 + x^1 e_2)(d\psi + \omega_1^2) + x^3(\omega_1^3 e_1 + \omega_2^3 e_2) + (x^1 \omega_1^3 + x^2 \omega_2^3) e_3 \\
&= -\left\{ \left(\sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} + \cosh \theta \cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} \right) du \right. \\
&\quad \left. + \left(\cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} + \cosh \theta \sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} \right) dv \right\} e \\
&\quad + \cosh \theta \left(\sin \frac{\phi - \phi_0}{2} du \cdot e_1 - \cos \frac{\phi - \phi_0}{2} dv \cdot e_2 \right) \\
&\quad + \sinh \theta \left(\sin \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} du \right. \\
&\quad \quad \left. - \cos \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} dv \right) e_3,
\end{aligned} \tag{3.13}$$

there we have

$$\begin{aligned}
 \xi_1^{*3} &= de_3^* \cdot e^* \\
 &= \sin \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} du + \cos \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} dv, \\
 \xi_2^{*3} &= de_3^* \cdot e^{*\perp} \\
 &= \sin \frac{\phi^* - \phi_0}{2} \sin \frac{\phi - \phi_0}{2} du - \cos \frac{\phi^* - \phi_0}{2} \cos \frac{\phi - \phi_0}{2} dv.
 \end{aligned} \tag{3.14}$$

Consider a local frame field $\{r^*; e_1^*, e_2^*, e_3^*\}$ on S^* such that

$$\begin{aligned}
 e^* &= \cos \frac{\phi - \phi_0}{2} e_1^* - \sin \frac{\phi - \phi_0}{2} e_2^*, \\
 e^{*\perp} &= \sin \frac{\phi - \phi_0}{2} e_1^* + \cos \frac{\phi - \phi_0}{2} e_2^*,
 \end{aligned} \tag{3.15}$$

and denote its dual as $\{\omega^{*1}, \omega^{*2}\}$. Then we have

$$\begin{aligned}
 \omega^{*1} &= \cos \frac{\phi - \phi_0}{2} \xi^{*1} + \sin \frac{\phi - \phi_0}{2} \xi^{*2} = -\frac{\lambda \sinh \gamma}{\sinh \theta} \cos \frac{\phi^*}{2} du, \\
 \omega^{*2} &= -\sin \frac{\phi - \phi_0}{2} \xi^{*1} + \cos \frac{\phi - \phi_0}{2} \xi^{*2} = -\frac{\lambda \sinh \gamma}{\sinh \theta} \sin \frac{\phi^*}{2} dv, \\
 \omega_1^{*3} &= \cos \frac{\phi - \phi_0}{2} \xi_1^{*3} + \sin \frac{\phi - \phi_0}{2} \xi_2^{*3} = \sin \frac{\phi^* - \phi_0}{2} du, \\
 \omega_2^{*3} &= -\sin \frac{\phi - \phi_0}{2} \xi_1^{*3} + \cos \frac{\phi - \phi_0}{2} \xi_2^{*3} = -\cos \frac{\phi^* - \phi_0}{2} dv.
 \end{aligned} \tag{3.16}$$

Comparing the above formulas with (3.1), we know that u and v are Tschebyscheff coordinates on S^* , and ϕ^* is its Tschebyscheff angle.

Remark 3.1. Theorem 3.1 implies a way to construct new Weingarten spacelike surfaces with condition (1.1) in R_1^3 from a given one. Without loss of generality, we can always assume $m \geq 0$ in (1.1) (if necessary, change the orientation of S).

THEOREM 3.2. Suppose S is a spacelike surface with condition (1.1) in R_1^3 , $m \geq 0$. For any given real number $\theta > 0$ and $\tanh(\theta/2) \neq m/\sqrt{m^2 + l^2}$, we can construct a Darboux line congruence Σ such that the solution of the completely integrable equation (3.4) is the Tschebyscheff angle of the corresponding surface S^* .

Proof. Assume that $\tanh(\theta/2) < m/\sqrt{m^2 + l^2}$. Then we can choose γ such that

$$\coth \gamma = \coth \frac{\theta}{2} \cdot \frac{m}{\sqrt{m^2 + l^2}} > 1.$$

Let (u, v) be the Tschebyscheff coordinates on S . Then Eq. (3.4) is completely integrable by Theorem 3.1 and Proposition 3.1. Let ϕ^* be the solution of (3.4) such that $\phi^*(u_0, v_0) = \phi(u_0, v_0)$. Put

$$\lambda = \frac{1}{\sqrt{m^2 + l^2}} \cdot \frac{\sinh \theta}{\sinh \gamma},$$

then (2.5), (2.11), (3.1) hold. Let

$$t = \sinh \gamma \left(\cos \frac{\phi^*}{2} \cdot e_1 - \sin \frac{\phi^*}{2} \cdot e_2 \right) + \cosh \gamma e_3, \quad (3.17)$$

and

$$r^* = r + \lambda t. \quad (3.18)$$

We want to prove that r^* is a spacelike surface, and that the above formula gives a timelike Darboux line congruence in R_1^3 associated with S and S^* .

Let

$$e_3^* = -\sinh \theta \left(\sin \frac{\phi^* - \phi_0}{2} e_1 + \cos \frac{\phi^* - \phi_0}{2} e_2 \right) - \cosh \theta e_3. \quad (3.19)$$

By differentiation of (3.18) we get

$$\begin{aligned} dr^* &= \omega^1 e_1 + \omega^2 e_2 - \lambda \sinh \gamma \left(\sin \frac{\phi^*}{2} e_1 + \cos \frac{\phi^*}{2} e_2 \right) \\ &\quad \times \left(\frac{\phi_u^* - \phi_v}{2} du + \frac{\phi_v^* - \phi_u}{2} dv \right) \\ &\quad + \lambda \cosh \gamma (\omega_1^3 e_1 + \omega_2^3 e_2) \\ &\quad + \lambda \sinh \gamma \left(\cos \frac{\phi^*}{2} \omega_1^3 - \sin \frac{\phi^*}{2} \omega_2^3 \right) e_3. \end{aligned} \quad (3.20)$$

By use of (3.1) and (3.4) we obtain

$$\begin{aligned}
 dr^* \cdot e_3^* &= -\sinh \theta \left(\sin \frac{\phi^* - \phi_0}{2} \omega^1 + \cos \frac{\phi^* - \phi_0}{2} \omega^2 \right) \\
 &\quad + \lambda \sinh \theta \sinh \gamma \cos \frac{\phi_0}{2} \left(\frac{\phi_u^* - \phi_v}{2} du + \frac{\phi_v^* - \phi_u}{2} dv \right) \\
 &\quad - \lambda \sinh \theta \cosh \gamma \left(\sin \frac{\phi^* - \phi_0}{2} \omega_1^3 + \cos \frac{\phi^* - \phi_0}{2} \omega_2^3 \right) \\
 &\quad + \lambda \cosh \theta \sinh \gamma \left(\cos \frac{\phi^*}{2} \omega_1^3 - \sin \frac{\phi^*}{2} \omega_2^3 \right) \\
 &= 0,
 \end{aligned} \tag{3.21}$$

thus e_3^* is a normal vector of S^* . From (3.19), $e_3^* \cdot e_3^* = -1$. Then S^* is a spacelike surface.

From the definition of t and e_3^* , we have

$$\begin{aligned}
 t \cdot e_3 &= -t \cdot e_3^* = -\cosh \gamma = \text{const.}, \\
 e_3 \cdot e_3^* &= \cosh \gamma = \text{const.}, \quad t \cdot t = -1,
 \end{aligned} \tag{3.22}$$

hence the line congruence given by (3.18) is a timelike Darboux line congruence. By Theorem 3.1, ϕ^* is the Tschebyscheff angle of S^* .

If $\tanh(\theta/2) > m/\sqrt{m^2 + l^2}$, then we can choose γ such that

$$\tanh \gamma = \coth \frac{\theta}{2} \cdot \frac{m}{\sqrt{m^2 + l^2}} < 1, \tag{3.23}$$

and let

$$t = \cosh \gamma \left(\cos \frac{\phi^*}{2} \cdot e_1 - \sin \frac{\phi^*}{2} \cdot e_2 \right) + \sinh \gamma e_3, \tag{3.24}$$

$$r^* = r + \lambda t. \tag{3.25}$$

The remainder of proof is similar to above; we omit the details here.

REFERENCES

1. W. H. Chen, Some results on spacelike surfaces in Minkowski 3-space, *Acta Math. Sinica* **37** (1994), 309–316. [In Chinese]
2. W. H. Chen and H. Z. Li, "Weingarten Surfaces and Sine-Gordon Equations," Research Report, No. 71, Inst. of Math. and School of Math. Scien., Peking University, 1996.

3. S. S. Chern and C. L. Terng, An analogue of Bäcklund's theorem in affine geometry, *Rocky Mountain J. Math.* **10** (1980), 105–124.
4. G. Darboux, "Lecons sur la Théorie Générale des Surfaces," Vol. 3, Gauthier-Villars, Paris, 1896.
5. L. P. Eisenhart, "A Treatise on the Differential Geometry of Curves and Surfaces," Dover, New York, 1960.
6. H. S. Hu, Soliton theory and differential geometry, in "Soliton Theory and Its Applications," pp. 343–395, Zhejiang Publishing House of Sci. and Tech., 1990.
7. Tian Chou, Bäcklund transformation on surfaces with $(k_1 - m)(k_2 - m) = -l^2$, ICTP Preprint, IC/95/296.
8. Y. Z. Huang, Bäcklund theorem on constant curvature surfaces in 3-dimensional Minkowski space and its higher dimensional generalization, *Acta Math. Sinica* **29** (1986), 684–690. [In Chinese]